

Flow at AGS Energies: A Barometer for High Density Effects?

D. E. Kahana¹, Y. Pang^{2,3}, E. Shuryak¹

¹*Physics Department, State University of New York,
Stony Brook, NY 11791, USA*

²*Physics Department, Brookhaven National Laboratory
Upton, NY 11973, USA*

³*Physics Department, Columbia University,
New York, NY 10027, USA*

(February 9, 2008)

Abstract

Preliminary data on transverse energy ‘flow’ and event asymmetries reported by the E877(814) collaborations are compared to ARC model calculations for Au+Au at full AGS beam energy. ARC triple differential cross-sections for protons and pions are presented. Proton flow is produced in ARC, with the maximum $\langle P_x \rangle \sim 120$ MeV/c. For central events $\langle P_x \rangle$ for the pions is near zero, consistent with experiment. Pion $\langle P_x \rangle$ opposite to the nucleons’ is evident in a peripheral sample, however, suggesting that this pion ‘anti-flow’ involves absorption by ‘spectator’ matter. ‘Squeeze-out’ of protons in central events at mid-rapidity is suggested by the ARC distributions.

It is presently of great interest to know how model calculations compare with data on flow at the AGS, especially since the excitation function of flow may well reveal new physics, if such is occurring at high baryon density and temperature [1,2]. ARC model calculations have previously been compared [1] to Bevalac data on ‘Sideways’ flow. Other aspects of flow, including the so-called longitudinal and radial flows, have also been extensively studied at Bevalac energies. Our study of Bevalac data was done at beam momenta in the range $p = 1\text{--}2$ GeV/c, for the heavy system Au+Au, energies an order of magnitude less than those for which the cascade was designed. Yet, very good agreement with observed flow was obtained, using a physically plausible treatment of two body scatterings.

Here we are concerned with recent measurements carried out at the AGS [3], by the E877(814) collaboration. For details of the experimental apparatus and other information we refer the reader to [4]. For our purposes it suffices to say that the instrument includes two highly segmented calorimeters, capable of measuring the energy deposition at various polar and azimuthal angles, as well as a forward spectrometer capable of particle identification. We will discuss calorimeter data on the azimuthal asymmetry of transverse energy as well as preliminary indications on the directed transverse momentum $\langle P_x \rangle$ measured in the spectrometer, using the reaction plane determined from the calorimeters.

The two calorimeters are TCAL and PCAL (target and participant calorimeters), with pseudorapidity coverages of $-0.5 < \eta < 0.8$ and $0.83 < \eta < 4.5$, respectively. Only PCAL is used in constructing the transverse energy distributions. The dispersion in the event plane determined from the PCAL is on the order of 60° (FWHM) from experiment [5], and theoretical simulation using ARC events yields a similar result. Therefore relatively large corrections to the raw experimental numbers are needed when extracting the triple differential cross-sections. Detector acceptances produce additional uncertainty. Nevertheless, the preliminary results clearly demonstrate the existence of proton flow at AGS, and give some hint as to the quantitative amount. For pions one observes, interestingly, no flow in a central sample [6].

Centrality cuts are placed on data and on ARC events using the total PCAL E_T . Central

events produce more transverse energy than peripheral. For ARC we calculated PCAL E_T using just the angular acceptance, not accounting for any efficiency corrections. The best procedure, of course, would be to pass ARC events through the experimental simulation of the PCAL response. However, the agreement in the E_T spectrum calculated directly is very close, certainly at better than 10% level for all but the most peripheral events. For the purpose of making centrality cuts on the flow measurements we have assumed that E877 E_T and (raw) ARC E_T are the same.

The azimuthal transverse energy distribution from E877 has been analysed [3], using an expansion in spherical harmonics. Moments v_n of the distribution of $E_T(\phi)$ are computed, by summing over the various cells $\{i\}$ composing the PCAL, which is nearly symmetric in ϕ :

$$v_n = \frac{\sum_i E_T^i \cos(n\phi_i)}{\sum_i E_T^i}$$

Moments up to v_4 have been considered. Including the corresponding sine transforms, one can define two-dimensional vectors v_n . The direction of v_1 for example, is an estimate of the direction of the reaction plane in a particular event. In principle then, the distributions of the relative phases of the v_n are meaningful too.

A more transparent procedure might have been to fit an ellipse to the distribution $E_T(\phi)$. The center, the major and minor axes, and the orientation of the ellipse would provide a small set of physically meaningful parameters by which to characterise the distributions. The finite detector segmentation in ϕ limits the highest frequency v_n containing significant information. In addition, fluctuations are large on an event by event basis, and high frequency noise is present, further obscuring the meaning of higher moments. We consider it unlikely that much real information is present beyond v_1 and v_2 , and it is hard to imagine what physics could lead to such azimuthal distributions. In any case, only v_1 shows any significant systematic effects (dependence on centrality for example) in the present data.

We have calculated v_1 and v_2 from ARC events, simulating the segmentation of PCAL using sixteen bins in ϕ . The pseudorapidity interval considered is $1.85 < \eta < 4.5$.

A comparison of the ARC moments with data is shown in Fig(1). One can see clearly that the distributions of v_n calculated from ARC agree quite well with the experimental ones, both in shape and in dependence on centrality. We plot only the magnitudes of v_n , and the Jacobian factor ($1/v_n$) is not included. The characters of the two dimensional distributions for v_1 and v_2 differ: the v_2 distributions have a maximum at $|v_2| = 0$ for all centralities. The distributions for v_1 are also peaked at zero for the most peripheral and most central event samples. But for semi-central events v_1 has a maximum *away* from zero. This is just the expected systematics of flow with centrality, the position of the maximum in v_1 being roughly related to the size of the transverse flow.

Plots of the triple differential cross sections from ARC, for an event sample corresponding to the E877 central cut (10% of PCAL E_t) are in Fig(2). For this cut, no flow is visible in the pion spectra. The pion distributions are symmetric around the beam axis, seemingly consistent with experiment [6]. The proton spectra shift progressively towards positive $\langle P_x \rangle$ with increasing rapidity. Additionally, at mid-rapidity the distribution is distorted, appearing roughly elliptical, with the major axis oriented along the $\langle P_y \rangle$ direction. This anisotropy vanishes near beam rapidity, where one sees only the shift toward positive $\langle P_x \rangle$. This would appear to be the so-called ‘squeeze out’, in which protons are ejected symmetrically out of the reaction plane near mid-rapidity. ‘Squeeze out’ has not been observed as yet at the AGS (it would be difficult at best to detect by analysing the v_n), but we note that quite a good definition of the reaction plane is probably necessary to discern it. Here, we are presenting essentially raw results from ARC. Folding these results with detector acceptances could possibly have a large effect.

For the peripheral event sample Fig (3) the triple differential cross-sections are quite different. Most strikingly, the pion distributions show a clear shift toward *negative* $\langle P_x \rangle$, *i. e.* opposite to the protons. This is the so-called ‘anti-flow’, and that it appears in the peripheral sample but is absent in the central one suggests that the effect is in large part due to shadowing by spectator nucleons. We note that E866 has detected an asymmetry in the pion distributions for semi-peripheral events [7].

Our previous study of heavy ion data at BEVALAC energies [1] using ARC has shown that a good description of the observed flow can be obtained across a range of energies and projectiles. Central to achieving this agreement at low energy was a correct treatment of pp scattering, specifically the collisions leading to two-body final states. Kinetic pressure in the cascade was found to produce on the order of 90% of the observed flow in the heaviest systems, once angular momentum conservation was properly taken into account by preserving the reaction plane in such collisions. If the reaction plane were randomized instead, about 70% of the flow would be obtained. Only an additional 10% difference in the flow is ascribed to a potential-like effect in two-body scattering [1]. Proton rapidity and transverse mass spectra were also found to be in agreement with data and to be nearly independent of the choice of orbits and the randomisation of the reaction plane. The conclusion is that at these low energies little potential or mean field enters: the pure cascade accounts for the flow.

Because the NN interaction is mostly repulsive at the higher LBL energies, the classical cascade can represent scattering well using dominantly repulsive outgoing paths for two-body final states. In actual potential modeling the character of the classical orbits will vary with impact parameter and energy. One expects repulsive orbits for small impact parameters probing the inner core, and attractive orbits passing through the outer regions. At sufficiently high energy the description in terms of real potential scattering breaks down, and pion production becomes the dominant process. A reasonable guess is that somewhere above the threshold for two pion production it is not sensible anymore to talk about potential models. In ARC, production of up to two pions $pp \rightarrow n_\pi \pi$ proceeds entirely through excitation of the $\Delta\Delta$ intermediate state. Hence this scattering can still be described in terms of two baryon states. For first collisions at AGS energies the average pion number has risen to 3.8 per inelastic collision, the two body optical theorem is violated and no description in terms of potential scattering is possible. The potential, if any, is clearly dominated by its absorptive, imaginary part, and the scattering has become diffractive in nature. What then can one say about the nature of the classical orbits at such high energies? Diffraction from a black disk occurs in the absence of either repulsion or attraction in the potential, therefore

the best guess is probably that the orbits should be equally often repulsive or attractive.

In constructing a cascade to be used at all energies between BEVALAC and AGS, one must interpolate between the mostly repulsive NN scattering occurring at low energies and the diffractive NN scattering taking place at higher energies. Heavy ion collisions at the AGS involve individual NN scatterings taking place at all energies equal to or below the first collision energy. The simplest procedure which builds in the essential physics is to use repulsive orbits below some energy, E_{min} , a 50/50 mixture of repulsive and attractive orbits above a higher energy, E_{max} , and to interpolate smoothly between these limits. In practice we take $E_{min} \sim 300$ MeV and $E_{max} \sim 500$ MeV corresponding crudely to the effective 2π and 3π thresholds. We can distinguish this approach quite sharply from one involving mean fields: what we propose is not at all a medium dependent effect, rather it is an energy dependent treatment of the *free space* two body scattering. This is justified, we think, by the nature of the NN interaction.

In Fig(4) we display the effect on $\langle P_x \rangle$ of choosing an energy dependent mixture of repulsive and attractive orbits versus the 50/50 mixture. Choosing purely repulsive orbits at BEVALAC energies produced a rather small ($< 10\%$) increase in flow discussed above. At AGS energies, however, the sensitivity to the classical orbits can be much larger, resulting in perhaps as much as 60% change in $\langle P_x \rangle$ for a central cut. Indeed, the ‘potential’ one would need to turn all the orbits outward for $p = 10.8$ GeV/c would have to be very large indeed and completely unrealistic. If all the orbits are made attractive, interestingly, the flow is reduced to near zero.

Summarising, present analysis of the very interesting E877 experiments on azimuthal asymmetry, combined with previous success at the BEVALAC, lends confidence to the cascade description of sideways flow. Certainly, the ARC calculations are in substantial agreement Fig.(1) with the higher energy data. The absolute flow magnitude is, however, more sensitive to the potential contribution at 11.6 GeV/c than at 2 GeV/c. The method of interpolation between real potential and diffractive two body scattering is straightforward, but its full validation awaits the detailed E877 flow results and triple differential cross-sections.

Nevertheless, it is interesting to speculate on the predictions of this model at intermediate beam momenta, $2 - 8$ GeV/c, relative to ongoing experiments with the EOS TPC [8], now installed at the AGS. At these, lower, AGS energies for $Au + Au$ one may reasonably expect to form a largely equilibrated compound system, still at high baryon density. So, the likelihood of nucleating appreciable amounts of quark-gluon plasma [9] should be optimized. The signature would be a considerable reduction in flow following from a reduction in kinetic pressure. The pioneering BEVALAC, E877 and E866 experiments provide needed guidance, fixing the flow at both low energy and at the full AGS Au energy, and setting the stage for perhaps dramatic results in between.

The authors thank S.H.Kahana for useful discussions. This work has been supported by DOE grants No. DE-FG02-93 ER40768, DE-AC02-76 CH00016, and DE-FG02-92 ER40699.

REFERENCES

- [1] D. E. Kahana *et al.*, Phys. Rev. Lett. **74**, 4404 (1995).
- [2] C. M. Hung, E. V. Shuryak, Phys. Rev. Lett. **75**, 4003 (1995). D. H. Rischke, M. Gyulassy, CU-TP-706, Sept. 1995.
- [3] J. Barrette *et al.*, Phys. Rev. Lett. **73**, 2532 (1994).
- [4] J. Simon-Gillo *et al.*, Nucl. Instrument. Methods **A 309**, 427 (1991).
- [5] Private Communication, P. Braun-Münzinger.
- [6] Private Communication, H. Wessels.
- [7] T. Abbott *et al.*, Phys. Rev. Lett. **70**, 1393 (1993).
- [8] G. Rai *et al.*, E895 (EOS) Collaboration.
- [9] J. Kapusta and R. Venugopalan, University of Minnesota Preprint, (1994)

FIGURES

Moments of Transverse Energy Distribution




FIG. 1. Moments of the distribution $E_t(\phi)$.

FIG. 2. Proton and pion transverse momentum distributions relative to the event plane (x-direction) for 10% central events. Pion distributions are symmetric at all rapidities, indicating no pion anti-flow at this centrality. This seems to be consistent with the current experimental picture. Proton distributions show a clear shift towards positive $\langle P_x \rangle$. A spectator component is present in the proton distribution for the most forward rapidity. The deformation of the proton distributions along $\langle P_y \rangle$ indicates proton squeezout occurs at mid-rapidity.

FIG. 3. Peripheral pion and proton momentum distributions ($b > 3$ fm). Proton distributions appear symmetric at all rapidities: $\langle P_x \rangle$ is small for peripheral samples. A large spectator component is present at the most forward rapidity. Pion distributions show a distinct shift to negative $\langle P_x \rangle$: that this pion ‘anti-flow’ appears only in the peripheral sample indicates it is largely due to shadowing.

Calculated Proton and Pion $\langle P_x \rangle$




FIG. 4. $\langle P_x \rangle$ for protons and pions. Central (10%) cut based on PCAL E_t . Energy dependent mixture of repulsive scattering against 50/50 mixture at all energies.

This figure "fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/nucl-th/9604008v1>

This figure "fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/nucl-th/9604008v1>